

循 環 器 内 科 学 英 文 原 著 論 文 実 績 (2010)

- 1) Abe H., **Takahashi M.**, Yaegashi H., Eda S., Tsunemoto H., Kamikozawa M., Koyama J., Yamazaki K., Ikeda U. : Efficacy of continuous positive airway pressure on arrhythmias in obstructive sleep apnea patients. *Heart Vessels* 25:63–69, 2010.
- 2) **Amagai Y.**, **Ishikawa S.**, Gotoh T., Kayaba K., Nakamura Y., **Kajii E.** : Sleep duration and incidence of cardiovascular events in a Japanese population: the Jichi Medical School cohort study. *J Epidemiol* 20:106–110, 2010.
- 3) Arakawa N., **Eguchi K.**, Ogawa T., Hashimoto R., Kato H. : Primary Aldosteronism Complicated by Severe Hypokalemic Myopathy. *Endocrinologist* 20:298–300, 2010.
- 4) Baba Y., **Ishikawa S.**, Kayaba K., Gotoh T., **Kajii E.** : High pulse pressure is associated with increased risk of stroke in Japanese: the JMS Cohort Study. *Blood Press* 20:10–14, 2010.
- 5) **Eguchi K.** : Tom Pickering as a clinical scientist: masked hypertension. *Blood Press Monit.* 15: 85–89, 2010.
- 6) **Eguchi K.**, **Hoshide S.**, **Hoshide Y.**, **Ishikawa S.**, **Shimada K.**, **Kario K.** : Reproducibility of ambulatory blood pressure in treated and untreated hypertensive patients. *J Hypertens* 28:918–924, 2010.
- 7) **Eguchi K.**, **Hoshide S.**, **Ishikawa S.**, **Shimada K.**, **Kario K.** : Short sleep duration is an independent predictor of stroke events in elderly hypertensive patients. *J Am Soc Hypertens* 4:255–262, 2010.
- 8) **Eguchi K.**, **Hoshide S.**, **Kabutoya T.**, **Shimada K.**, **Kario K.** : Is very low dose hydrochlorothiazide combined with candesartan effective in uncontrolled hypertensive

patients? Blood Press Monit 15:308–311, 2010.

9) **Eguchi K., Kario K.**: Is nondipping harmful in normotensive, healthy subjects? Am J Hypertens 23:222–223, 2010.

10) **Eguchi K., Matsui Y., Shibasaki S., Hoshide S., Kabutoya T., Ishikawa J., Ishikawa S., Shimada K., Kario K.**: Controlling evening BP as well as morning BP is important in hypertensive patients with prediabetes/diabetes: the JMS-1 study. Am J Hypertens 23:522–527, 2010.

11) **Eguchi K., Schwartz JE., Pickering TG., Hoshide S., Ishikawa J., Shimada K., Kario K.**: Increased heart rate variability during sleep is a predictor for future cardiovascular events in patients with type 2 diabetes. Hypertens Res;33:737–742, 2010.

12) **Fukutomi M., Kario K.**: Aging and hypertension. Expert Rev Cardiovasc Ther 8:1531–1539, 2010.

13) Hasegawa T., Boden-Albala B., **Eguchi K.**, Jin Z., Sacco RL., Homma S., Di Tullio MR.: Impaired flow-mediated vasodilatation is associated with increased left ventricular mass in a multiethnic population. The Northern Manhattan Study. Am J Hypertens 23:413–419, 2010.

14) Hasegawa Y., **Shimada K.**, Yamaguchi T.: Efficacy and safety of diuretics in combination with perindopril in hypertensive stroke patients: Results of the Japan Perindopril and Diuretics on Cerebrovascular Disease study (J-PADOC). Blood Press Suppl 19:10–16, 2010.

15) Hirashima C., Ohkuchi A., Takahashi K., Suzuki H., Yoshida M., Ohmaru T., **Eguchi K.**, Ariga H., Matsubara S., Suzuki M.: Gestational hypertension as a subclinical

preeclampsia in view of serum levels of angiogenesis-related factors. *Hypertens Res* 34:212–217, 2010.

- 16) Hirose H., **Ishikawa S.**, Gotoh T., **Kabutoya T.**, Kayaba K., **Kajii E.**: Cardiac mortality of premature ventricular complexes in healthy people in japan. *J Cardiol* 56:23–26, 2010.
- 17) **Ichida M.**, **Katsurada K.**, **Komori T.**, **Matsumoto J.**, **Ohkuchi A.**, **Izumi A.**, **Matsubara S.**, **Mitsuhashi T.**, **Kario K.**: Effectiveness of bromocriptine treatment in a patient with peripartum cardiomyopathy. *J Cardiol Case* 2:e28–e31, 2010.

18) Inoue H., Nozawa T., Hirai T., Goto S., Origasa H., **Shimada K.**, Uchiyama S., Hirabayashi T., Koretsune Y., Ono S., Hasegawa T., Sasagawa Y., Kaneko Y., Ikeda Y.: Sex-related differences in the risk factor profile and medications of patients with atrial fibrillation recruited in J-TRACE. *Circ J* 74:650–654, 2010.

- 19) Ise H., Kobayashi S., Goto M., Sato T., Kawakubo M., **Takahashi M.**, Ikeda U., Akaike T.: Vimentin and desmin possess GlcNAc-binding lectin-like properties on cell surfaces. *Glycobiology* 20:843–864.

20) **Ishikawa J.**, Haimoto H., **Hoshide S.**, **Eguchi K.**, **Shimada K.**, **Kario K.**, **J-HOP Study Investigators Group.**: An increased visceral–subcutaneous adipose tissue ratio is associated with difficult-to-treat hypertension in men. *J Hypertens* 28:1340–1346, 2010.

- 21) **Ishikawa J.**, **Hoshide S.**, **Eguchi K.**, **Schwartz JE.**, **Pickering TG.**, **Shimada K.**, **Kario K.**: Masked hypertension defined by ambulatory blood pressure monitoring is associated with an increased serum glucose level and urinary albumin–creatinine ratio. *J Clin Hypertens (Greenwich)* 12:578–587, 2010.

- 22) **Ishikawa Y., Ishikawa J., Ishikawa S., Kajii E., Schwartz JE., Pickering TG., Kario K., Jichi Medical School Cohort Investigators Group.**: Prehypertension and the risk for cardiovascular disease in the Japanese general population: The Jichi Medical School Cohort Study. *J Hypertens* 28:1630–1637, 2010.
- 23) Ishiwata A., **Mimuro J., Mizukami H., Kashiwakura Y., Yasumoto A., Sakata A., Ohmori T., Madoiwa S., Ono F., Shima M., Yoshioka A., Ozawa K., Sakata Y.**: Mutant macaque factor IX T262A: a tool for hemophilia B gene therapy studies in macaques. *Thromb Res* 125: 533–537, 2010.
- 24) **Kabutoya T., Hoshide S., Ishikawa J., Eguchi K., Shimada K., Kario K.**: The effect of pulse rate and blood pressure dipping status on the risk of stroke and cardiovascular disease in Japanese hypertensive patients. *Am J Hypertens* 23 :749–755, 2010.
- 25) **Kabutoya T., Ishikawa J., Hoshide S., Eguchi K., Shimada K., Kario K.**: Poor blood pressure and urinary albumin excretion responses to home blood pressure-based antihypertensive therapy in depressive hypertensive patients. *J Clin Hypertens (Greenwich)* 12:345–349, 2010.
- 26) **Kabutoya T., Mitsuhashi T., Hata Y., Hashimoto T., Nakagami R., Osada J., Watanabe T., Shimada K., Kario K.**: Beneficial effects of upgrading from right ventricular pacing to cardiac resynchronization therapy in patients with heart failure compared to de novo cardiac resynchronization therapy. *J Arrhythmia* 26: 16–20, 2010.
- 27) **Kario K.**: Thomas G. Pickering – a great mentor. *Blood Press Monit* 15:82–84, 2010.

- 28) **Kario K.** : Measuring the effects of stress on the cardiovascular system during a disaster: the effective use of self-measured blood pressure monitoring. *J Hypertens* 28:657–659, 2010.
- 29) **Kario K.** : Proposal of RAS-diuretic vs. RAS-calcium antagonist strategies in high-risk hypertension: insight from the 24-hour ambulatory blood pressure profile and central pressure. *J Am Soc Hypertens* 4:215–218, 2010.
- 30) **Kario K.** : Morning surge in blood pressure and cardiovascular risk: evidence and perspectives. *Hypertension* 56:765–773, 2010.
- 31) **Kario K., Hoshide S., Shimizu M., Yano Y., Eguchi K., Ishikawa J., Ishikawa S., Shimada K.** : Effect of dosing time of angiotensin II receptor blockade titrated by self-measured blood pressure recordings on cardiorenal protection in hypertensives: the Japan Morning Surge–Target Organ Protection (J-TOP) study. *J Hypertens* 28:1574–1583, 2010.
- 32) **Kario K., Shirayama M., Hiramatsu K., Shiosakai K., Sugiyama M., Shimada K.** : A new baroreceptor sensitivity-restoring Ca-channel blocker diminishes age-related morning blood pressure increase in hypertensive patients: open-label monitoring of azelnidipine treatment for hypertension in the early morning (At-HOME) Study. *Pharmaceuticals* 3: 225–236, 2010.
- 33) **Kario K., Yano Y., Matsuo T., Hoshide S., Eguchi K., Shimada K.** : Additional impact of morning haemostatic risk factors and morning blood pressure surge on stroke risk in older Japanese hypertensive patients. *Eur Heart J* 32:574–580, 2010.
- 34) **Kimura M., Murakami T., Kizaka-Kondoh S., Itoh M., Yamamoto K., Hojo Y., Takano**

M., **Kario K.**, **Shimada K.**, Kobayashi E.: Functional molecular imaging of ILK-mediated Akt/PKB signaling cascades and the associated role of beta-parvin. *J Cell Sci* 123:747–755, 2010.

35) **Kondo H.**, **Hojo Y.**, **Takahashi N.**, **Ikemoto T.**, **Aoki H.**, **Dezaki K.**, **Kario K.**, **Katsuki T.**, Yada T., **Shimada K.**: Roles of ghrelin in left-ventricular remodelling

after acute myocardial infarction. *Heart Asia* 2: 1–4, 2010.

36) Kumazaki S., Koyama J., Aizawa K., Kasai H., Koshikawa M., Izawa A., Tomita T., **Takahashi M.**, Ikeda U.: Effect of graft adaptation of the internal mammary artery on

longitudinal phasic blood flow velocity characteristics after surgery. *Heart Vessels* 25:515–521, 2010.

37) **Matsui Y.**, **Eguchi K.**, **Ishikawa J.**, **Shimada K.**, **Kario K.**: Urinary albumin

excretion during angiotensin II receptor blockade: comparison of combination

treatment with a diuretic or a calcium-channel blocker. *Am J Hypertens* Epub 2010 Dec 16.

38) **Matsui Y.**, **Eguchi K.**, O'Rourke MF., **Ishikawa J.**, **Shimada K.**, **Kario K.**:

Association between aldosterone induced by antihypertensive medication and arterial stiffness reduction: The J-CORE study. *Atherosclerosis* 215:184–188, 2010.

39) **Matsui Y.**, **Eguchi K.**, **Shibasaki S.**, **Ishikawa J.**, **Hoshide S.**, **Shimada K.**, **Kario K.**:

Impact of arterial stiffness reduction on urinary albumin excretion during antihypertensive treatment: The Japan morning surge-1 study. *J Hypertens* 28:1752–1760, 2010.

40) **Matsui Y.**, **Eguchi K.**, **Shibasaki S.**, **Ishikawa J.**, **Shimada K.**, **Kario K.**: Morning

hypertension assessed by home monitoring is a strong predictor of concentric left ventricular hypertrophy in patients with untreated hypertension. *J Clin Hypertens (Greenwich)* 12:776–783, 2010.

41) Matsumoto M., **Ishikawa S., Kajii E.**: Rurality of communities and incidence of stroke: A confounding effect of weather conditions? *Rural Remote Health* 10:1493, 2010.

42) Matsumoto M., **Ishikawa S., Kajii E.**: Cumulative effects of weather on stroke incidence: a multi-community cohort study in Japan. *J Epidemiol* 20:136–142, 2010.

43) **Miyashita H.**, Aizawa A., Hashimoto J., Hirooka Y., Imai Y., Kawano Y., Kohara K., Sunagawa K., Suzuki H., Tabara Y., Takazawa K., Takenaka T., Yasuda H., **Shimada K.** : Cross-sectional characterization of all classes of antihypertensives in terms of central blood pressure in Japanese hypertensive patients. *Am J Hypertens* 23:260–268, 2010.

44) **Nagai M., Hoshide S., Kario K.** : Hypertension and dementia. *Am J Hypertens* 23:116–124, 2010.

45) **Nagai M., Hoshide S., Kario K.** : Sleep duration as a risk factor for cardiovascular disease- a review of the recent literature. *Curr Cardiol Rev* 6:54–61, 2010.

46) **Nagai M., Hoshide S., Kario .** : The insular cortex and cardiovascular system: a new insight into the brain-heart axis. *J Am Soc Hypertens* 4:174–182, 2010.

47) Naganuma M., Koga M., Shiokawa Y., Nakagawara J., Furui E., Kimura K., Yamagami H., Okada Y., Hasegawa Y., **Kario K.**, Okuda S., Nishiyama K., Minematsu K., Toyoda K.: Reduced estimated glomerular filtration rate is associated with stroke outcome after

intravenous rt-PA: the Stroke Acute Management with Urgent Risk-Factor Assessment and Improvement (SAMURAI) rt-PA registry. *Cerebrovasc Dis* 31:123–129, 2010.

48) Nezu T., Koga M., Kimura K., Shiokawa Y., Nakagawara J., Furui E., Yamagami H., Okada Y., Hasegawa Y., **Kario K.**, Okuda S., Nishiyama K., Naganuma M., Minematsu K., Toyoda K.: Pretreatment ASPECTS on DWI predicts 3-month outcome following rt-PA: SAMURAI rt-PA Registry. *Neurology* 75:555–561, 2010.

49) Niwa Y., **Ishikawa S.**, Gotoh T., Kayaba K., Nakamura Y., **Kajii E.**: Association between stroke and metabolic syndrome in a Japanese population: Jichi Medical School (JMS) Cohort Study. *J Epidemiol* 20:62–69, 2010.

50) Parati G., Ochoa JE., Ramos C., **Hoshide S.**, Lonati L., Bilo G.: Efficacy and tolerability of olmesartan/amlodipine combination therapy in patients with mild-to-severe hypertension: focus on 24-h blood pressure control. *Ther Adv Cardiovasc Dis* 4:301–313, 2010.

51) Parati G., Stergiou GS., Asmar R., Bilo G., de Leeuw P., Imai Y., **Kario K.**, Lurbe E., Manolis A., Mengden T., O'Brien E., Ohkubo T., Padfield P., Palatini P., **Pickering TG.**, Redon J., Revera M., Ruilope LM., Shennan A., Staessen JA., Tisler A., Waeber B., Zanchetti A., Mancia G.: European Society of Hypertension practice guidelines for home blood pressure monitoring. *J Hum Hypertens* 24:779–785, 2010.

52) **Shimada K.**: Will the "JMU System" Stop Medical System Collapse? : *JMAJ* 53: 1–2, 2010.

53) **Shimizu M.**, **Kario K.**: Perfect 24-Hour BP Management for High-Risk Hypertension.

大韓高血圧学会誌 ; 16 、 2010 .

- 54) **Takahashi M.** : Immature erythroid cells: a new source for therapeutic angiogenesis? J Mol Cell Cardiol 49:341–342, 2010.
- 55) Teramoto T., **Shimada K.**, Uchiyama S., Sugawara M., Goto Y., Yamada N., Oikawa S., Ando K., Ishizuka N., Yamazaki T., Yokoyama K., Murata M., Ikeda Y.: Rationale, design, and baseline data of the Japanese Primary Prevention Project (JPPP)-a randomized, open-label, controlled trial of aspirin versus no aspirin in patients with multiple risk factors for vascular events. Am Heart J 159:361–369 e364, 2010.
- 56) Uchiyama S., Shibata Y., Hirabayashi T., Mihara B., Hamashige N., Kitagawa K., Goto S., Origasa H., **Shimada K.**, Kobayashi H., Isozaki M., Ikeda Y.: Risk factor profiles of stroke, myocardial infarction, and atrial fibrillation: a Japanese Multicenter Cooperative Registry. J Stroke Cerebrovasc Dis 19:190–197, 2010.
- 57) Uurtuya S., **Kotani K.**, **Koibuchi H.**, **Miyamoto M.**, **Kario K.**, **Yamada T.**, **Taniguchi N.** : The relationship between serum C-reactive protein and daily physical activity in Japanese hypertensive patients. Clin Exp Hypertens 32:517–522, 2010.
- 58) Uurtuya S., **Kotani K.**, **Taniguchi N.**, Yoshioka H., **Kario K.**, **Ishibashi S.**, **Yamada T.**, **Kawano M.**, Khurelbaatar N., **Itoh K.**, Lkhagvasuren T.: Comparative study of atherosclerotic parameters in Mongolian and Japanese patients with hypertension and diabetes mellitus. J Atheroscler Thromb 17:181–188, 2010.
- 59) **Yano Y.**, **Hoshide S.**, **Tamaki N.**, Nagata M., Sasaki K., Kanemura Y., **Shimada K.**, **Kario K.** : Efficacy of Eplerenone added to Renin-Angiotensin Blockade in Elderly Hypertensive Patients –The Jichi –Eplerenone Treatment (JET) study– . J Renin Angiotensin Aldosteron Syst :2010.

- 60) **Yano Y.**, Inokuchi T., **Hoshide S.**, Kanemaru Y., **Shimada K.**, **Kario K.**: Association of poor physical function and cognitive dysfunction with high nocturnal blood pressure level in treated elderly hypertensive patients. *Am J Hypertens* 24:285-291, 2010.
- 61) **Yano Y.**, **Kario K.**: Therapeutic implications of high-dose angiotensin receptor blocker monotherapy in mild-to-moderate hypertensive patients. *Hypertens Res* 33:981-983, 2010.
- 62) **Yano Y.**, Matsuda S., Hatakeyama K., Sato Y., Imamura T., **Shimada K.**, Kodama T., **Kario K.**, Asada Y.: Plasma pentraxin 3, but not high-sensitivity C-reactive protein, is a useful inflammatory biomarker for predicting cognitive impairment in elderly hypertensive patients. *J Gerontol A Biol Sci Med Sci* 65:547-552, 2010.
- 63) 長田直人、成尾浩明、江川久子、**矢野裕一朗**、白尾英仁、松田俊太郎、金丸勝弘：末梢血管抵抗の増減に対する橈骨動脈の脈波増大係数（AI）の変動について。日本臨床生理学会雑誌 40巻2号；83-87.
- 64) **吉村裕子**、外山卓二：慢性腎臓病の有無は初発急性心筋梗塞患者の遠隔期主要心血管イベント発症に寄与しうる。群馬県核医学研究会雑誌 1.24, No2;38-40.