

循環器内科学英文原著論文実績（2011）

- 1) Eguchi K.: Ambulatory blood pressure monitoring in diabetes and obesity-a review. *Int J Hypertens*;2:954757,2011.
- 2) Eguchi K.: A call for wider use of ambulatory blood pressure monitoring in patients with diabetes. *Hypertens Res*;34:1171-1172,2011.
- 3) Eguchi K., Kuruvilla S., Ishikawa J., Ogedegbe G., Gerin W., Schwartz JE., Pickering TG.: Correlations between different measures of clinic, home, and ambulatory blood pressure in hypertensive patients. *Blood Press Monit*;16:142-148,2011.
- 4) Eguchi K., Tsujita A., Ono S., Kario K.: A case of syncope induced in the supine position. *Int J Hypertens*; 342140,2011.
- 5) Fukutomi M., Hoshide S., Eguchi K., Watanabe T., Shimada K., Kario K.: Differential effects of strict blood pressure lowering by losartan/hydrochlorothiazide combination therapy and high-dose amlodipine monotherapy on microalbuminuria: The ALPHABET study. *J Am Soc Hypertens*;6:73-82,2011.
- 6) Hata T., Takahashi M., Hida S., Kawaguchi M., Kashima Y., Usui F., Morimoto H., Nishiyama A., Izawa A., Koyama J., Iwakura Y., Taki S., Ikeda U.: Critical role of Th17 cells in inflammation and neovascularization after ischaemia. *Cardiovasc Res*;90:364-372,2011.
- 7) Hoshide S., Parati G., Matsui Y., Shibasaki S., Eguchi K., Kario K.: Orthostatic hypertension: Home blood pressure monitoring for detection and assessment of treatment with doxazosin. *Hypertens Res*;35:100-106,2011.
- 8) Ikemoto T., Hojo Y., Kondo H., Takahashi N., Hirose M., Nishimura Y., Katsuki T., Shimada K., Kario K.: Plasma endoglin as a marker to predict cardiovascular events in patients with chronic coronary artery diseases. *Heart Vessels* ,2011.
- 9) Ise M., Ise H., Shiba Y., Kobayashi S., Goto M., Takahashi M., Akaike T., Ikeda U.: Targeting N-acetylglucosamine-bearing polymer-coated liposomes to vascular smooth muscle cells. *J Artif Organs*;14:301-309,2011.
- 10) Ishikawa J., Eguchi K., Hoshide S., Shimada K., Kario K.: Relationship between the change in left ventricular hypertrophy and asleep blood pressure after sodium restriction and/or diuretic treatment. *Blood Press Monit*;16:172-179,2011.

- 11) Ishikawa J., Ishikawa Y., Edmondson D., Pickering T., Schwartz JE: Age and the difference between awake ambulatory blood pressure and office blood pressure: A meta-analysis. *Blood Press Monit*;16:159-167,2011.
- 12) Izumi M., Iwata T., Ichida M., Sakano Y., Konishi H., Kurumizawa S., Tanakanashi S., Shimada M., Kario K.: Left Ventricular Apical and Atrial Appendage Thrombus with Hypertrophic Cardiomyopathy in Sinus Rhythm , a case report.:J of Medical Ultrasonics;online first ;2011.
- 13) Izumi M., Suzuki K., Sakamoto T., Hayashi M.: Advantages and limitations of antihypertensive treatment for stroke risk in a general population: the Akita Stroke Registry. *Acta Cardiol*;66:729-735,2011.
- 14) Izumi M., Teraoka S., Yamashita K., Matsumoto K., Muronoi T., Izawa Y., Yonekura C., Ano M and Suzukawa M.: Successful Management of Aortic Thrombi Resulted in Spinal Cord infarction in a Patient with Antiphospholipid Antibody Syndrome(APS)during Hospital Course of Acute Cholecystitis. *Int Medical Case Reports J*;4:93-96,2011.
- 15) Kabutoya T., Mitsuhashi T., Watanabe T., Nakagami R., Hata Y., Shimada K., Kario K.: The relationship between optimization for cardiac resynchronization therapy by measurement of dp/dt and the middle-to-long-term prognosis of heart failure patients.:J of Arrythmia;27:208-213,2011.
- 16) Kario K.: Are melatonin and its receptor agonist specific antihypertensive modulators of resistant hypertension caused by disrupted circadian rhythm? *J Am Soc Hypertens*;5:354-358, 2011.
- 17) Kario K., Nishizawa M., Hoshide S., Shimpou M., Ishibashi Y., Kunii O., Shibuya K.: Development of a disaster cardiovascular prevention network. *Lancet*;378:1125-1127,2011.
- 18) Kario K., Yano Y., Matsuo T., Hoshide S., Asada Y., Shimada K.: Morning blood pressure surge, morning platelet aggregation, and silent cerebral infarction in older Japanese hypertensive patients. *J Hypertens*;29:2433-2439,2011.
- 19) Kario K., Yano Y., Matsuo T., Hoshide S., Eguchi K., Shimada K.: Additional impact of morning haemostatic risk factors and morning blood pressure surge on stroke risk in older Japanese hypertensive patients. *Eur Heart J*;32:574-580,2011.
- 20) Kawaguchi M., Takahashi M., Hata T., Kashima Y., Usui F., Morimoto H., Izawa A., Takahashi Y., Masumoto J., Koyama J., Hongo M., Noda T., Nakayama J., Sagara J., Taniguchi S., Ikeda U.: Inflammasome activation of cardiac fibroblasts is essential for myocardial ischemia/reperfusion injury. *Circulation*;123:594-604,2011.

- 21) Kawano H., Yamamoto H., Miyata S., Izumi M., Hirano T., Toratani N., Kakutani I., Sheppard JA., Warkentin TE., Kada A., Sato , Okamoto S., Nagatsuka K., Naritomi H., Toyoda K., Uchino M., Minematsu K.: Prospective multicentre cohort study of heparin-induced thrombocytopenia in acute ischaemic stroke patients. *Br J Haematol*;154:378-386,2011.
- 22) Kenzaka T., Shimoshikiryō M., Kitao A., Kario K., Hashimoto M.: Positron emission tomography scan can be a reassuring tool to treat difficult cases of infective endocarditis. *J Nucl Cardiol*;18:741-743,2011.
- 23) Kinugawa S., Tojo A., Sakai T., Tsumura H., Takahashi M., Hirata Y., Fujita T.: Mechanism of selective albuminuria in puromycin nephritic rats and its blockade by a NADPH oxidase inhibitor, *Kidney Int*;80:1328-1338,2011.
- 24) Koga M., Kimura K., Shibasaki K., Shiokawa Y., Nakagawara J., Furui E., Yamagami H., Okada Y., Hasegawa Y., Kario K., Okuda S., Naganuma M., Nezu T., Maeda K., Minematsu K., Toyoda K.: CHADS2 score is associated with 3-month clinical outcomes after intravenous rt-PA therapy in stroke patients with atrial fibrillation: SAMURAI rt-PA registry. *J Neurol Sci*;306:49-53, 2011.
- 25) Koga M., Shiokawa Y., Nakagawara J., Furui E., Kimura K., Yamagami H., Okada Y., Hasegawa Y., Kario K., Okuda S., Endo K., Miyagi T., Osaki M., Minematsu K., Toyoda K.: Low-dose intravenous recombinant tissue-type plasminogen activator therapy for patients with stroke outside European indications: Stroke acute management with urgent Risk-factor assessment and improvement (SAMURAI) rt - PA registry. *Stroke*;43:253-255,2011.
- 26) Kotani K., Yamada T., Miyamoto M., Kario K., Ishibashi S., Taniguchi N.: The correlation between the cardio-ankle vascular index (CAVI) and serum amyloid A in asymptomatic Japanese subjects. *Heart Vessels* ,2011
- 27) Maeda K., Koga M., Okada Y., Kimura K., Yamagami H., Okuda S., Hasegawa Y., Shiokawa Y., Furui E., Nakagawara J., Kario K., Nezu T., Minematsu K., Toyoda K.: Nationwide survey of neuro-specialists' opinions on anticoagulant therapy after intracerebral hemorrhage in patients with atrial fibrillation. *J Neurol Sci*;312:82-85,2011.
- 28) Masuda T., Murata M., Honma S., Iwazu Y., Sasaki N., Ogura M., Onishi A., Ando Y., Muto S., Shimada K., Kario K., Kusano E., Asano Y.: Sleep-disordered breathing predicts cardiovascular events and mortality in hemodialysis patients. *Nephrol Dial Transplant*;26:2289-2295,2011.
- 29) Matsui Y., Eguchi K., Ishikawa J., Shimada K., Kario K.: Urinary albumin excretion during

- angiotensin II receptor blockade: Comparison of combination treatment with a diuretic or a calcium-channel blocker. *Am J Hypertens*;24:466-473,2011.
- 30) Matsui Y., Eguchi K., O'Rourke MF., Ishikawa J., Shimada K., Kario K.: Association between aldosterone induced by antihypertensive medication and arterial stiffness reduction: The J-CORE study. *Atherosclerosis*;215:184-188,2011.
- 31) Matsui Y., Ishikawa J., Eguchi K., Shibasaki S., Shimada K., Kario K: Maximum value of home blood pressure: A novel indicator of target organ damage in hypertension. *Hypertension*;57:1087-1093,2011.
- 32) Matsui Y., Ishikawa J., Shibasaki S., Shimada K., Kario K.: Association between home arterial stiffness index and target organ damage in hypertension: Comparison with pulse wave velocity and augmentation index. *Atherosclerosis*;219:637-642,2011.
- 33) Matsuoka H., Ando K., Ueshima K., Babazono T., Kario K., Tanaka S., Nakao K., Fujita T.: Design and rationale of the study of assessment for kidney function by urinary microalbumin in randomized (SAKURA) trial. *Clin Exp Hypertens*;33:455-462,2011.
- 34) Motoki H., Koyama J., Tomita T., Aizawa K., Kasai H., Koshikawa M., Izawa A., Kumazaki S., Takahashi M., Ikeda U.: Transient pseudorestrictive pattern of transmural flow velocity curve in patients with paroxysmal atrial fibrillation. *Echocardiology*;28:289-297,2011.
- 35) Nagai M., Hoshide S., Ishikawa J., Shimada K., Kario K.: Visit-to-visit blood pressure variations: New independent determinants for carotid artery measures in the elderly at high risk of cardiovascular disease. *J Am Soc Hypertens*;5:184-192.
- 36) Nagai M., Hoshide S., Kario K.: Visit-to-visit blood pressure variability and carotid artery atherosclerosis: Heart rate was not a confounder. *Hypertension*;58:e16,2011.
- 37) Nagai M., Kario K.: Chronic kidney disease, 24-h blood pressure burden and their effects on silent cerebral injury and cognitive impairment: Might age serve as a modulator? *Hypertens Res*;34:1253-1254,2011
- 38) Naganuma M., Koga M., Shiokawa Y., Nakagawara J., Furui E., Kimura K., Yamagami H., Okada Y., Hasegawa Y., Kario K., Okuda S., Nishiyama K., Minematsu K., Toyoda K.: Reduced estimated glomerular filtration rate is associated with stroke outcome after intravenous rt-PA: The stroke acute management with urgent risk-factor assessment and improvement (SAMURAI) rt-PA registry. *Cerebrovasc Dis*;31:123-129,2011.

- 39) Nezu T., Koga M., Nakagawara J., Shiokawa Y., Yamagami H., Furui E., Kimura K., Hasegawa Y., Okada Y., Okuda S., Kario K., Naganuma M., Maeda K., Minematsu K., Toyoda K.: Early ischemic change on CT versus diffusion-weighted imaging for patients with stroke receiving intravenous recombinant tissue-type plasminogen activator therapy: Stroke acute management with urgent risk-factor assessment and improvement (SAMURAI) rt-PA registry. *Stroke*;42:2196-2200, 2011.
- 40) Nishikawa M., Shimpo M., Hashimoto T., Murata M., Shimada K., Kario K.: A case of infected aortic aneurysm with possible intramural abscess resolved through discharge into the vascular lumen without surgical intervention. *J Cardiol Cases* : 3 ; e163-e166, 2011.
- 41) Ohmori T., Yano Y., Sakata A., Ikemoto T., Shimpo M., Madoiwa S., Katsuki T., Mimuro J., Shimada K., Kario K., Sakata Y.: Lack of association between serum paraoxonase-1 activity and residual platelet aggregation during dual antiplatelet therapy. *Thromb Res*.Nov.23 ,2011.
- 42) Sakata T., Kario K.: Antiplatelet therapy effectively reduces plasma plasminogen activator inhibitor-1 levels. *Atherosclerosis*;214:490-491,2011.
- 43) Saito Y., Kondo H., Hojo Y.: Granzyme B as a novel factor involved in cardiovascular disease.*J Cardiol*.:57(2),141-147,2011.
- 44) Sekine H., Shimizu T., Dobashi I., Matsuura K., Hagiwara N., Takahashi M., Kobayashi E., Yamato M., Okano T.: Cardiac cell sheet transplantation improves damaged heart function via superior cell survival in comparison with dissociated cell injection. *Tissue Eng Part A*;17:2973-2980,2011.
- 45) Shimada K., Kario K.: Preface - 12th Japanese annual conference on chronocardiology and hypertension. *Clin Exp Hypertens*;33:209,2011.
- 46) Shimizu M., Ishikawa J., Yano Y., Hoshide S., Shimada K., Kario K.: The relationship between the morning blood pressure surge and low-grade inflammation on silent cerebral infarct and clinical stroke events. *Atherosclerosis*;219:316-321,2011.
- 47) Shirasaki O., Kuwabara M., Saito M., Tagami K., Washiya S., Kario K.: Development and clinical application of a new technique for detecting 'sleep blood pressure surges' in sleep apnea patients based on a variable desaturation threshold. *Hypertens Res*;34:922-928,2011.
- 48) Suzuki K., Izumi M., Sakamoto T., Hayashi M.: Blood pressure and total cholesterol level are critical risks especially for hemorrhagic stroke in Akita, Japan. *Cerebrovasc Dis*;31: 100-106,2011.

- 49) Takahashi S., Ito T., Zenimaru Y., Suzuki J., Miyamori I., Takahashi M., Ishida T., Hirata K., Yamamoto T., Iwasaki T., Hattori H., Shiomi M.: Species differences of macrophage very low-density-lipoprotein (VLDL) receptor protein expression. *Biochem Biophys Res Commun*; 407:656-662.
- 50) Umegaki H., Haimoto , Hoshide S., Kario K.: Strong association between insulin resistance with urinary albumin excretion in older adults. *J Am Geriatr Soc*;59:1361-1362.
- 51) Vongpatanasin W., Kario K., Atlas SA., Victor RG.: Central sympatholytic drugs. *J Clin Hypertens (Greenwich)*;13:658-661,2011.
- 52) Wang JG., Kario K., Lau T., Wei YQ., Park CG., Kim CH., Huang J., Zhang W., Li Y., Yan P., Hu D.: Use of dihydropyridine calcium channel blockers in the management of hypertension in Eastern Asians: A scientific statement from the Asian Pacific Heart Association. *Hypertens Res*;34:423-430,2011.
- 53) Yamaguchi A.,Murakami T.,Takahashi M.,Kobayashi E.,Sugawara Y.: Luminescence imaging of regenerating free bone graft in rats. *Plast Reconstr Surg*;127:78-87,2011.
- 54) Yano Y., Haimoto H., Hoshide S., Kabutoya T., Eguchi K., Kario K.: Evening heart rate measured at home is associated with visceral obesity and abnormal fat distribution in patients with hypertension. *Am J Hypertens*;24:783-788,2011.
- 55) Yano Y., Hoshide S., Etoh T., Tamaki N., Yokota N., Kario K.: Synergistic effect of chronic kidney disease and high circulatory norepinephrine level on stroke risk in Japanese hypertensive patients. *Atherosclerosis*;219:273-279,2011.
- 56) Yano Y., Hoshide S., Shimizu M., Eguchi K., Ishikawa J., Ishikawa S., Shimada K., Kario K.: Association of Home and Ambulatory Blood Pressure changes with changes in cardiovascular biomarkers during antihypertensive treatment. *Am J Hypertens*;25:306-312,2011.
- 57) Yano Y., Hoshide S., Tamaki N., Inokuchi T., Nagata M., Yokota N., Hidaka T., Kanemaru Y., Matsuda S., Kuwabara M., Shimada K., Kario K.: Regional differences in hypertensive cardiovascular remodeling between fishing and farming communities in Japan. *Am J Hypertens*;24:437-443,2011.
- 58) Yano Y., Hoshide S., Tamaki N., Nagata M., Sasaki K., Kanemaru Y., Shimada K., Kario K.: Efficacy of eplerenone added to renin-angiotensin blockade in elderly hypertensive patients: The Jichi-Eplerenone Treatment (JET) study. *J Renin Angiotensin Aldosterone Syst*;12:340-347,2011.

- 59) Yano Y., Inokuchi T., Hoshide S., Kanemaru Y., Shimada K., Kario K.: Association of poor physical function and cognitive dysfunction with high nocturnal blood pressure level in treated elderly hypertensive patients. Am J Hypertens;24:285-291,2011.
- 60) Yano Y., Kario K.: The risk of cold temperature: An important aspect of the determination of morning blood pressure surge. Hypertens Res;34:36-38,2011.
- 61) 江口和男 : ミニ・レビュー 圧受容体反射感受性と高血圧性臓器障害の関連性。自律神経、5:p363-365, 2011。
- 62) 桂田健一, 上野修市, 宮原義典, 横田克明, 簿 義仁, 新保昌久, 三橋武司, 河田政明, 莢尾七臣 : 心房中隔欠損による重症肺動脈性肺高血圧症の治療方針。Therapeutic Research、1264-66、2011。
- 63) 甲谷友幸、三橋武司、渡部智紀、中神理恵子、簿 義仁、島田和幸、筈尾七臣:冠静脈内で逆行性の遅伝導路の焼灼に成功した三重伝導路をもつ slow-slow 型房室結節リエントリ頻拍の 1 例。心電図 : 31、 p 249-255、 2011。