

2013 年

[81]Yano Y, Kario K, Ishikawa S, Ojima T, Gotoh T, Kayaba K, et al. Associations between diabetes, leanness, and the risk of death in the Japanese general population: the Jichi Medical School Cohort Study. *Diabetes Care*. 2013 May;36(5):1186-92.

[80] Takezako Y, Ishikawa S, Kajii E. Advance directives in Japanese nursing homes. *J Pain Symptom Manage*. 2013 Jan;45(1):63-70 e7.

[79] Kaneda Y, Ishikawa S, Sadakane A, Gotoh T, Kayaba K, Yasuda Y, et al. Insulin Resistance and Risk of Cerebral Infarction in a Japanese General Population: The Jichi Medical School Cohort Study. *Asia Pac J Public Health*. 2013 Mar 27;doi: 10.1177/1010539513480233.

[78] Kaneda Y, Ishikawa S, Sadakane A, Gotoh T, Kayaba K, Yasuda Y, et al. Fasting insulin and risk of cerebral infarction in a Japanese general population: The Jichi Medical School Cohort Study. *Neurology Asia*. 2013;18(4):343-8.

[77] Kaneda Y, Ishikawa S, Gotoh T, Kayaba K, Yasuda Y, Kajii E. Risk of Cerebral Infarction assessed by Fasting Insulin and Fasting Glucose in a Japanese General Population: The Jichi Medical School Cohort Study. *Jichi Medical University Journal*. 2013;36:in press.

[76] Kakehi E, Kotani K, Ishikawa S, Gotoh T, Kayaba K, Nakamura Y, et al. Serum Non-High-Density Lipoprotein Cholesterol Levels and the Incidence of Ischemic Stroke in a Japanese Population: The Jichi Medical School Cohort Study. *Asia Pac J Public Health*. 2013 Feb 20: doi: 10.1177/1010539513475649

[75] Ishikawa S, Kotani K, Kario K, Kayaba K, Gotoh T, Nakamura Y, et al. Inverse association between serum lipoprotein(a) and cerebral hemorrhage in the Japanese population. *Thromb Res*. 2013 Feb;131(2):e54-8.