[19] Tsutsumi A, Kayaba K, Ishikawa S, Kario K, Matsuo H, Takuma S. [Jichi Medical School Social Support Scale (JMS-SSS) revision and tests for validity and reliability]. Nippon Koshu Eisei Zasshi. 2000 Oct;47(10):866-78.

[18] Kayaba K, Ishikawa S, Gotoh T, Nago N, Kajii E, Nakamura Y, et al. Five-year intra-individual variability in C-reactive protein levels in a Japanese population-based study: the Jichi Medical School Cohort Study at Yamato, 1993-1998. Jpn Circ J. 2000 Apr;64(4):303-8.

[17] Ishikawa S, Deguchi T, Hara K, Takuma S, Kayaba K, Tsutsumi A, et al. Lipoprotein(a) levels and apolipoprotein(a) isoforms related to life style risk factors. J Epidemiol. 1999 Feb;9(1):32-9.

- [16] Tsutsumi A, Tsutsumi K, Kayaba K, Theorell T, Nago N, Kario K, et al. Job strain and biological coronary risk factors: a cross-sectional study of male and female workers in a Japanese rural district. Int J Behav Med. 1998;5(4):295-311.
- [15] Tsutsumi A, Tsutsumi K, Kayaba K, Igarashi M. Health-related behaviors, social support, and community morale. Int J Behav Med. 1998;5(2):166-82.
- [14] Kayaba K, Nago N, Miyamoto T, Mizooka M, Terada M, Kario K, et al. Glycated hemoglobin levels and their correlation with atherosclerotic risk factors in a Japanese population--the Jichi Medical School Cohort Study 1993-1995. Jpn Circ J. 1998 Apr;62(4):261-6.
- [13] Kayaba K, Igarashi M, Okamoto H, Tsuda F. Prevalence of anti-hepatitis C antibodies in a rural community without high mortality from liver disease in Niigata prefecture. J Epidemiol. 1998 Oct;8(4):250-5.
- [12] Ishikawa S, Goto T, Nago N. Effect of moderate alcohol consumption on Lp(a) lipoprotein concentrations. Reduction is not found in women. Bmj. 1998 May 30;316(7145):1675-6.

[11] Ishikawa S, Kario K, Nago N, Kayaba K, Hiraoka J, Matsuo H, et al. Factor VII and fibrinogen levels examined by age, sex, and other atherosclerotic risk factors in a Japanese population. The Jichi Medical School Cohort Study. Thromb Haemost. 1997 May;77(5):890-3.

[10] Inoue K, Nago N, Matsuo H, Goto T, Miyamoto T, Saegusa T, et al. Serum insulin and lipoprotein(a) concentrations. The Jichi Medical School Cohort Study. Diabetes Care. 1997 Aug;20(8):1242-7.

[9] Tsutsumi K, Tsutsumi A, Orth-Gomer K. The effect of labor force participation on coronary heart disease risk factors among middle-aged women: a cross-sectional study in a Japanese rural district. Int J Behav Med. 1996;3(4):370-86.

[8]Nishio H, Lee MJ, Fujii M, Kario K, Kayaba K, Shimada K, et al. A common mutation in methylenetetrahydrofolate reductase gene among the Japanese population. Jpn J Hum Genet. 1996 Jun;41(2):247-51.

[7]Kario K, Nago N, Kayaba K, Saegusa T, Matsuo H, Goto T, et al. Characteristics of the insulin resistance syndrome in a Japanese population. The Jichi Medical School Cohort Study. Arterioscler Thromb Vasc Biol. 1996 Feb;16(2):269-74.

[6]Kario K, Kanai N, Saito K, Nago N, Matsuo T, Shimada K. Ischemic stroke and the gene for angiotensin-converting enzyme in Japanese hypertensives. Circulation. 1996 May 1;93(9):1630-3.

- [5] Nishiuma S, Kario K, Kayaba K, Nagio N, Shimada K, Matsuo T, et al. Effect of the angiotensinogen gene Met235-->Thr variant on blood pressure and other cardiovascular risk factors in two Japanese populations. J Hypertens. 1995 Jul;13(7):717-22.
- [4] Nago N, Kayaba K, Hiraoka J, Matsuo H, Gotoh T, Kario K, et al. Lipoprotein(a) levels in the Japanese population: influence of age and sex, and relation to atherosclerotic risk factors: the Jichi Medical School Cohort Study. Am J Epidemiol. 1995;141(9):815-21.
- [3] Gotoh T, Kuroda T, Yamasawa M, Nishinaga M, Mitsuhashi T, Seino Y, et al. Correlation between lipoprotein(a) and aortic valve sclerosis assessed by echocardiography (the JMS Cardiac Echo and Cohort Study). Am J Cardiol. 1995 Nov 1;76(12):928-32.

[2]Kario K, Matsuo T, Imiya M, Kayaba K, Kuroda T, Nago N, et al. Close relation between lipoprotein (a) levels and atherothrombotic disease in Japanese subjects > 75 years of age. Am J Cardiol. 1994 Jun 15;73(16):1187-90.

[1]Hiraoka J, Nakamura Y, Yanagawa H, Nago N. Distribution of Lipoprotein (a) and relationships between its level and blood chemical findings in a rural area in Japan. J Epidemiol. 1994;4(3):163-39.