

2008 年

[45] Matsumoto M, Ishikawa S, Kajii E. Adiponectin and noncardiovascular death: a nested case-control study. *Metabolism*. 2008;57(6):811-8.

[44] Matsumoto M, Ishikawa S, Kajii E. Association of adiponectin with cerebrovascular disease: a nested case-control study. *Stroke*. 2008 Feb;39(2):323-8.

[43] Ishikawa Y, Ishikawa J, Ishikawa S, Kayaba K, Nakamura Y, Shimada K, et al. Prevalence and determinants of prehypertension in a Japanese general population: the Jichi Medical School Cohort Study. *Hypertension Research*. 2008;31(7):1323-30.

[42] Ishikawa S, Kayaba K, Gotoh T, Nago N, Nakamura Y, Tsutsumi A, et al. Incidence of total stroke, stroke subtypes, and myocardial infarction in the Japanese population: the JMS Cohort Study. *J Epidemiol*. 2008;18(4):144-50.

[41] Ishikawa S, Kario K, Kayaba K, Gotoh T, Nago N, Nakamura Y, et al. Continued high risk of stroke in treated hypertensives in a general population: the Jichi Medical School Cohort study. *Hypertens Res*. 2008 Jun;31(6):1125-33.

[40] Hirokawa K, Tsutsumi A, Kayaba K. Psychosocial job characteristics and plasma fibrinogen in Japanese male and female workers: the Jichi Medical School cohort study. *Atherosclerosis*. 2008 Jun;198(2):468-76.