

2009 年

[53] Tsutsumi A, Kayaba K, Kario K, Ishikawa S. Prospective study on occupational stress and risk of stroke. *Arch Intern Med.* 2009 Jan 12;169(1):56-61.

[52] Sadakane A, Gotoh T, Ishikawa S, Nakamura Y, Kayaba K. Amount and frequency of alcohol consumption and all-cause mortality in a Japanese population: the JMS Cohort Study. *J Epidemiol.* 2009;19(3):107-15.

[51] Matsumoto M, Ishikawa S, Kayaba K, Gotoh T, Nago N, Tsutsumi A, et al. Risk charts illustrating the 10-year risk of myocardial infarction among residents of Japanese rural communities: the JMS Cohort Study. *J Epidemiol.* 2009;19(2):94-100.

[50] Ishikawa S, Matsumoto M, Kayaba K, Gotoh T, Nago N, Tsutsumi A, et al. Risk charts illustrating the 10-year risk of stroke among residents of Japanese rural communities: the JMS Cohort Study. *J Epidemiol.* 2009;19(2):101-6.

[49] Ishikawa J, Ishikawa S, Kabutoya T, Gotoh T, Kayaba K, Schwartz JE, et al. Cornell product left ventricular hypertrophy in electrocardiogram and the risk of stroke in a general population. *Hypertension.* 2009 Jan;53(1):28-34.

[48] Hirokawa K, Tsutsumi A, Kayaba K. Occupation and plasma fibrinogen in Japanese male and female workers: The Jichi Medical School Cohort study. *Soc Sci Med.* 2009 Jan 23.

[47] Hayasaka S, Shibata Y, Ishikawa S, Kayaba K, Gotoh T, Noda T, et al. Physical activity and all-cause mortality in Japan: the Jichi Medical School (JMS) Cohort Study. *J Epidemiol.* 2009;19(1):24-7.

[46] Hatano Y, Matsumoto M, Ishikawa S, Kajii E. Plasma adiponectin level and myocardial infarction: the JMS Cohort Study. *J Epidemiol.* 2009;19(2):49-55.