

內科學英文原著論文業績

1. Akimoto T, Saito O, Inoue M, Nishino K, Onishi A, Kotoda A, Ando Y, Muto S, Kusano E. Rapid formation of Aspergillus mycetoma in a patient receiving corticosteroid treatment. Serial radiographic observation over two months. *Intern Med* 2007;46:733-7.
2. Aoki H, Ohnishi H, Hama K, Shinozaki S, Kita H, Osawa H, Yamamoto H, Sato K, Tamada K, Sugano K. Cyclooxygenase-2 is required for activated pancreatic stellate cells to respond to proinflammatory cytokines. *Am J Physiol Cell Physiol* 2007;292:C259-68.
3. Bailie GR, Elder SJ, Mason NA, Asano Y, Cruz JM, Fukuhara S, Lopes AA, Mapes DL, Mendelsohn DC, Bommer J, Young EW. Sexual dysfunction in dialysis patients treated with antihypertensive or antidepressive medications: results from the DOPPS. *Nephrol Dial Transplant* 2007;22:1163-70.
4. Choi YL, Kaneda R, Wada T, Fujiwara S, Soda M, Watanabe H, Kurashina K, Hatanaka H, Enomoto M, Takada S, Yamashita Y, Mano H. Identification of a constitutively active mutant of JAK3 by retroviral expression screening. *Leuk Res* 2007;31:203-9.
5. Dalmau J, Tuzun E, Wu HY, Masjuan J, Rossi JE, Voloschin A, Baehring JM, Shimazaki H, Koide R, King D, Mason W, Sansing LH, Dichter MA, Rosenfeld MR, Lynch DR. Paraneoplastic anti-N-methyl-D-aspartate receptor encephalitis associated with ovarian teratoma. *Ann Neurol* 2007;61:25-36.
6. Eguchi K, Boden-Albala B, Jin Z, Di Tullio MR, Rundek T, Rodriguez CJ, Homma S, Sacco RL. Usefulness of fasting blood glucose to predict vascular outcomes among individuals without diabetes mellitus (from the Northern Manhattan Study). *Am J Cardiol* 2007;100:1404-9.
7. Eguchi K, Ishikawa J, Hoshide S, Ishikawa S, Pickering TG, Schwartz JE, Homma S, Shimada K, Kario K. Differential impact of left ventricular mass and relative wall thickness on cardiovascular prognosis in diabetic and nondiabetic hypertensive subjects. *Am Heart J* 2007;154:79 e9-15.
8. Eguchi K, Ishikawa J, Hoshide S, Pickering TG, Shimada K, Kario K. Masked hypertension in diabetes mellitus: a potential risk. *J Clin Hypertens (Greenwich)* 2007;9:601-7.
9. Eguchi K, Kario K. The morning-evening difference in self-measured blood pressure: a potential predictor of cardiovascular disease. *Hypertens Res* 2007;30:877-8.
10. Eguchi K, Matsui Y, Shibasaki S, Ishikawa J, Hoshide S, Ishikawa S, Kabutoya T, Schwartz JE, Pickering TG, Shimada K, Kario K. Age-specific impact of self-monitored pulse pressure on hypertensive target organ damage in treated hypertensive patients. *J Clin Hypertens (Greenwich)* 2007;9:522-9.
11. Eguchi K, Matsui Y, Shibasaki S, Ishikawa J, Hoshide S, Ishikawa S, Kabutoya T, Schwartz JE, Pickering TG, Shimada K, Kario K. Changes in self-monitored pulse pressure correlate with improvements in B-type natriuretic Peptide and urinary albumin in treated hypertensive patients. *Am J Hypertens* 2007;20:1268-75.
12. Eguchi K, Pickering TG, Kario K. Why is blood pressure so hard to control in patients

with type 2 diabetes? *J Cardiometab Syndr* 2007;2:114-8.

13. Eguchi K, Schwartz JE, Roman MJ, Devereux RB, Gerin W, Pickering TG. Metabolic syndrome less strongly associated with target organ damage than syndrome components in a healthy, working population. *J Clin Hypertens (Greenwich)* 2007;9:337-44.
14. Eguchi K, Tomizawa H, Ishikawa J, Hoshide S, Fukuda T, Numao T, Shimada K, Kario K. Effects of new calcium channel blocker, azelnidipine, and amlodipine on baroreflex sensitivity and ambulatory blood pressure. *J Cardiovasc Pharmacol* 2007;49:394-400.
15. Eguchi K, Tomizawa H, Ishikawa J, Hoshide S, Numao T, Fukuda T, Shimada K, Kario K. Comparison of the effects of pioglitazone and metformin on insulin resistance and hormonal markers in patients with impaired glucose tolerance and early diabetes. *Hypertens Res* 2007;30:23-30.
16. Eguchi K, Tomizawa H, Ishikawa J, Hoshide S, Pickering TG, Shimada K, Kario K. Factors associated with baroreflex sensitivity: association with morning blood pressure. *Hypertens Res* 2007;30:723-8.
17. Eguchi K, Yacoub M, Jhalani J, Gerin W, Schwartz JE, Pickering TG. Consistency of blood pressure differences between the left and right arms. *Arch Intern Med* 2007;167:388-93.
18. Ethier J, Bragg-Gresham JL, Piera L, Akizawa T, Asano Y, Mason N, Gillespie BW, Young EW. Aspirin prescription and outcomes in hemodialysis patients: the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Am J Kidney Dis* 2007;50:602-11.
19. Fujiwara S-I, Muroi K, Kikuchi S, Kawano-Yamamoto C, Matsuyama T, Mori M, Nagai T, Akutsu M, Ozawa K. Development of streptococcus meningitis and Epstein-Barr virus reactivation after non-T-cell-depleted human leukocyte antigen-haploidential peripheral blood stem cell transplantation based on feto-maternal microchimerism. *Leuk Lymphoma* 2007;48:640-2.
20. Fujiwara S, Yamashita Y, Choi YL, Watanabe H, Kurashina K, Soda M, Enomoto M, Hatanaka H, Takada S, Ozawa K, Mano H. Transforming activity of purinergic receptor P2Y, G protein coupled, 8 revealed by retroviral expression screening. *Leuk Lymphoma* 2007;48:978-86.
21. Fukuda H, Sonoo M, Ishibashi M. Muscle afferent contributions to tibial nerve somatosensory evoked potentials investigated using knee stimulations. *Clin Neurophysiol* 2007;118:2104-11.
22. Fukuda H, Sonoo M, Kako M, Shimizu T. Optimal method to determine the stimulus intensity for median nerve somatosensory evoked potentials. *J Clin Neurophysiol* 2007;24:358-62.
23. Fukumoto A, Tanaka S, Yamamoto H, Yao T, Matsui T, Iida M, Goto H, Sakamoto C, Chiba T, Sugano K. Diagnosis and treatment of small-bowel stricture by double balloon endoscopy. *Gastrointest Endosc* 2007;66:S108-12.
24. Gao J, Katagiri H, Ishigaki Y, Yamada T, Ogihara T, Imai J, Uno K, Hasegawa Y, Kanzaki M, Yamamoto TT, Ishibashi S, Oka Y. Involvement of apolipoprotein E in excess fat accumulation and insulin resistance. *Diabetes* 2007;56:24-33.
25. Hatanaka H, Takada S, Choi YL, Fujiwara S, Soda M, Enomoto M, Kurashina K,

- Watanabe H, Yamashita Y, Sugano K, Mano H. Transforming activity of purinergic receptor P2Y, G-protein coupled, 2 revealed by retroviral expression screening. *Biochem Biophys Res Commun* 2007;356:723-6.
26. Hayashino Y, Fukuhara S, Akiba T, Akizawa T, Asano Y, Saito A, Bragg-Gresham JL, Ramirez SP, Port FK, Kurokawa K. Diabetes, glycaemic control and mortality risk in patients on haemodialysis: the Japan Dialysis Outcomes and Practice Pattern Study. *Diabetologia* 2007;50:1170-7.
27. Hirahara I, Inoue M, Okuda K, Ando Y, Muto S, Kusano E. The potential of matrix metalloproteinase-2 as a marker of peritoneal injury, increased solute transport, or progression to encapsulating peritoneal sclerosis during peritoneal dialysis--a multicentre study in Japan. *Nephrol Dial Transplant* 2007;22:560-7.
28. Hirohata S, Arinuma Y, Takayama M, Yoshio T. Association of cerebrospinal fluid anti-ribosomal p protein antibodies with diffuse psychiatric/neuropsychological syndromes in systemic lupus erythematosus. *Arthritis Res Ther* 2007;9:R44.
29. Hoshide S, Ishikawa J, Eguchi K, Ojima T, Shimada K, Kario K. Masked nocturnal hypertension and target organ damage in hypertensives with well-controlled self-measured home blood pressure. *Hypertens Res* 2007;30:143-9.
30. Ideno J, Mizukami H, Kakehashi A, Saito Y, Okada T, Urabe M, Kume A, Kuroki M, Kawakami M, Ishibashi S, Ozawa K. Prevention of diabetic retinopathy by intraocular soluble flt-1 gene transfer in a spontaneously diabetic rat model. *Int J Mol Med* 2007;19:75-9.
31. Imai E, Horio M, Iseki K, Yamagata K, Watanabe T, Hara S, Ura N, Kiyohara Y, Hirakata H, Moriyama T, Ando Y, Nitta K, Inaguma D, Narita I, Iso H, Wakai K, Yasuda Y, Tsukamoto Y, Ito S, Makino H, Hishida A, Matsuo S. Prevalence of chronic kidney disease (CKD) in the Japanese general population predicted by the MDRD equation modified by a Japanese coefficient. *Clin Exp Nephrol* 2007;11:156-63.
32. Imai E, Horio M, Nitta K, Yamagata K, Iseki K, Hara S, Ura N, Kiyohara Y, Hirakata H, Watanabe T, Moriyama T, Ando Y, Inaguma D, Narita I, Iso H, Wakai K, Yasuda Y, Tsukamoto Y, Ito S, Makino H, Hishida A, Matsuo S. Estimation of glomerular filtration rate by the MDRD study equation modified for Japanese patients with chronic kidney disease. *Clin Exp Nephrol* 2007;11:41-50.
33. Ishibashi S. The vascular renin-angiotensin system as a possible source of vascular inflammation in fructose-fed rats. *Hypertens Res* 2007;30:375-6.
34. Ishikawa J, Kario K. Assessment of nocturnal blood pressure by home blood pressure monitoring. *Hypertens Res* 2007;30:661-2.
35. Ishikawa J, Kario K. Alcohol consumption reduces coronary heart disease even among men with hypertension. *Expert Review of Cardiovascular Therapy* 2007;633-634 .
36. Ishikawa J, Tamura Y, Hoshide S, Eguchi K, Ishikawa S, Shimada K, Kario K. Low-grade inflammation is a risk factor for clinical stroke events in addition to silent cerebral infarcts in Japanese older hypertensives: the Jichi Medical School ABPM Study, wave 1. *Stroke* 2007;38:911-7.
37. Ishikawa S, Kayaba K, Gotoh T, Nakamura Y, Kario K, Ito Y, Kajii E. Comparison of

- C-reactive protein levels between serum and plasma samples on long-term frozen storage after a 13.8 year interval: the JMS Cohort Study. *J Epidemiol* 2007;17:120-4.
38. Ishikawa S, Kazuomi K, Kayaba K, Gotoh T, Nago N, Nakamura Y, Tsutsumi A, Kajii E. Linear relationship between blood pressure and stroke: the Jichi Medical School Cohort Study. *J Clin Hypertens (Greenwich)* 2007;9:677-83.
39. Ishikawa S, Shibano Y, Asai Y, Kario K, Kayaba K, Kajii E. Blood pressure categories and cardiovascular risk factors in Japan: the Jichi Medical School (JMS) Cohort Study. *Hypertens Res* 2007;30:643-9.
40. Ishikawa T, Morita M, Nakano I. Constant blood flow reduction in premotor frontal lobe regions in ALS with dementia - a SPECT study with 3D-SSP. *Acta Neurol Scand* 2007;116:340-4.
41. Ito T, Okada T, Mimuro J, Miyashita H, Uchibori R, Urabe M, Mizukami H, Kume A, Takahashi M, Ikeda U, Sakata Y, Shimada K, Ozawa K. Adenoassociated virus-mediated prostacyclin synthase expression prevents pulmonary arterial hypertension in rats. *Hypertension* 2007;50:531-6.
42. Ito T, Okada T, Miyashita H, Nomoto T, Nonaka-Sarukawa M, Uchibori R, Maeda Y, Urabe M, Mizukami H, Kume A, Takahashi M, Ikeda U, Shimada K, Ozawa K. Interleukin-10 expression mediated by an adeno-associated virus vector prevents monocrotaline-induced pulmonary arterial hypertension in rats. *Circ Res* 2007;101:734-41.
43. Ito T, Ozawa K, Shimada K. Current drug targets and future therapy of pulmonary arterial hypertension. *Curr Med Chem* 2007;14:719-33.
44. Iwamoto M, Hayashi Y, Yamamoto H, Kita H, Yano T, Sunada K, Sugano K, Horie H, Nagai H, Kanai N. Double-balloon endoscopy for jejunal hemangioma (with video). *Gastrointest Endosc* 2007;66:184-5; discussion 185.
45. Iwazu Y, Honma S, Fujisawa G, Uki K, Yanaka I, Sato Y, Murata M, Kusano E, Asano Y. Hyponatremic seizure associated with acute respiratory infection. *Clin Exp Nephrol* 2007;11:230-4.
46. Izumiya M, Tamada K, Tomiyama T, Hanatsuka K, Ohashi A, Sugano K. A case of choledochocoele mimicking choledocholithiasis on MR cholangiography: A new differential diagnosis of pseudo-filling defect. *J Magn Reson Imaging* 2007;26:1097-100.
47. Jiang T, Wang XX, Scherzer P, Wilson P, Tallman J, Takahashi H, Li J, Iwahashi M, Sutherland E, Arend L, Levi M. Farnesoid X receptor modulates renal lipid metabolism, fibrosis, and diabetic nephropathy. *Diabetes* 2007;56:2485-93.
48. Kamata Y, Iwamoto M, Minota S. Consecutive use of sildenafil and bosentan for the treatment of pulmonary arterial hypertension associated with collagen vascular disease: sildenafil as reliever and bosentan as controller. *Lupus* 2007;16:901-3.
49. Kamata Y, Takahashi Y, Iwamoto M, Matsui K, Murakami Y, Muroi K, Ikeda U, Shimada K, Yoshio T, Okazaki H, Minota S. Local implantation of autologous mononuclear cells from bone marrow and peripheral blood for treatment of ischaemic digits in patients with connective tissue diseases. *Rheumatology (Oxford)* 2007;46:882-4.
50. Kario K. Vascular damage in exaggerated morning surge in blood pressure.

Hypertension 2007;49:771-2.

51. Kario K. Nondipping in nocturnal blood pressure in diabetes: an indicator of autonomic nervous dysfunction? *Am J Hypertens* 2007;20:546-7.
52. Kario K. Preceding linkage between a morning surge in blood pressure and small artery remodeling: an indicator of prehypertension? *J Hypertens* 2007;25:1573-5.
53. Kikuchi J, Shimizu R, Wada T, Ando H, Nakamura M, Ozawa K, Furukawa Y. E2F-6 suppresses growth-associated apoptosis of human hematopoietic progenitor cells by counteracting proapoptotic activity of E2F-1. *Stem Cells* 2007;25:2439-47.
54. Kikuchi S, Nagai T, Kunitama M, Kiritto K, Ozawa K, Komatsu N. Active FKHLR1 overcomes imatinib resistance in chronic myelogenous leukemia-derived cell lines via the production of tumor necrosis factor-related apoptosis-inducing ligand. *Cancer Sci* 2007;98:1949-58.
55. Kimata N, Albert JM, Akiba T, Yamazaki S, Kawaguchi Y, Fukuhara S, Akizawa T, Saito A, Asano Y, Kurokawa K, Pisoni RL, Port FK. Association of mineral metabolism factors with all-cause and cardiovascular mortality in hemodialysis patients: the Japan dialysis outcomes and practice patterns study. *Hemodial Int* 2007;11:340-8.
56. Kita H, Yamamoto H. New indications of double balloon endoscopy. *Gastrointest Endosc* 2007;66:S57-9.
57. Kita H, Yamamoto H, Yano T, Miyata T, Iwamoto M, Sunada K, Arashiro M, Hayashi Y, Ido K, Sugano K. Double balloon endoscopy in two hundred fifty cases for the diagnosis and treatment of small intestinal disorders. *Inflammopharmacology* 2007;15:74-7.
58. Kitatani N, Taniguchi A, Fukushima M, Nakai Y, Nagasaka S, Ohgushi M, Ohya M, Kuroe A, Inagaki N, Seino Y. Association of serum resistin with TNF system activity in Japanese type 2 diabetic patients. *Diabetes Metab* 2007;33:156-7.
59. Kobayashi M, Okada T, Murakami T, Ozawa K, Kobayashi E, Morita T. Tissue-targeted in vivo gene transfer coupled with histone deacetylase inhibitor depsipeptide (FK228) enhances adenoviral infection in rat renal cancer allograft model systems. *Urology* 2007;70:1230-6.
60. Kraemer FB, Shen WJ, Patel S, Osuga J, Ishibashi S, Azhar S. The LDL receptor is not necessary for acute adrenal steroidogenesis in mouse adrenocortical cells. *Am J Physiol Endocrinol Metab* 2007;292:E408-12.
61. Laflamme MA, Chen KY, Naumova AV, Muskheli V, Fugate JA, Dupras SK, Reinecke H, Xu C, Hassanipour M, Police S, O'Sullivan C, Collins L, Chen Y, Minami E, Gill EA, Ueno S, Yuan C, Gold J, Murry CE. Cardiomyocytes derived from human embryonic stem cells in pro-survival factors enhance function of infarcted rat hearts. *Nat Biotechnol* 2007;25:1015-24.
62. Madoiwa S, Someya T, Hironaka M, Kobayashi H, Ohmori T, Mimuro J, Sugiyama Y, Morita T, Nishimura Y, Tarumoto T, Ozawa K, Saito K, Sakata Y. Annexin 2 and hemorrhagic disorder in vascular intimal carcinomatosis. *Thromb Res* 2007;119:229-40.
63. Matsui Y, Eguchi K, Ishikawa J, Hoshide S, Shimada K, Kario K. Subclinical arterial damage in untreated masked hypertensive subjects detected by home blood pressure measurement. *Am J Hypertens* 2007;20:385-91.

64. Mensink PB, Haringsma J, Kucharzik T, Cellier C, Perez-Cuadrado E, Monkemuller K, Gasbarrini A, Kaffles AJ, Nakamura K, Yen HH, Yamamoto H. Complications of double balloon enteroscopy: a multicenter survey. *Endoscopy* 2007;39:613-5.
65. Miyoshi T, Nagai T, Kikuchi S, Ohmine K, Nakamura M, Hanafusa T, Komatsu N, Ozawa K. Cloning and characterization of a human BCR/ABL-positive cell line, K562/RR, resistant to the farnesyltransferase inhibition by tipifarnib. *Exp Hematol* 2007;35:1358-65.
66. Mori M, Furukawa Y, Takatoku M, Nagai T, Muroi K, Ozawa K. B-cell lymphoma showing typical features of angioimmunoblastic T-cell lymphoma. *Jichi Med Univ J* 2007;30:89-95.
67. Morimatsu Y, Matsubara S, Hirose N, Ohkuchi A, Izumi A, Ozaki K, Ozawa K, Suzuki M. Acute promyelocytic leukemia: an unusual cause showing prolonged disseminated intravascular coagulation after placental abruption. *Arch Gynecol Obstet* 2007;277:e267-270.
68. Muramatsu S. Acupuncture and knee osteoarthritis. *Ann Intern Med* 2007;146:147; author reply 148-9.
69. Muroi R, Kusaka I, Kawakami A, Rokkaku K, Nagasaka S, Ishibashi S. Thyrotropin receptor antibodies measured by second-generation assay are highly specific for Graves' disease and correlate with serum thyrotropin levels in patients with Graves' disease in remission and during maintenance therapy. *Jichi Medical University Journal* 2007;30:19-28.
70. Mutoh H, Hayakawa H, Sakamoto H, Sugano K. Homeobox protein CDX2 reduces Cox-2 transcription by inactivating the DNA-binding capacity of nuclear factor-kappaB. *J Gastroenterol* 2007;42:719-29.
71. Nabeya D, Taniguchi A, Fukushima M, Nakai Y, Matsumoto K, Kuroe A, Ohgushi M, Ohya M, Nagasaka S, Inagaki N, Seino Y. Soluble vascular cell adhesion molecule-1 is independently associated with soluble tumor necrosis factor receptor 2 in Japanese type 2 diabetic patients. *Diabetes Metab* 2007;33:81-2.
72. Nagasaka S, Sato N, Takahashi N, Kusaka I, Ishibashi S. New insights on the simultaneous assessment of insulin sensitivity and beta-cell function with the HOMA2 method: response to Caumo et al. *Diabetes Care* 2007;30:e42; author reply e43.
73. Nakata M, Okada T, Ozawa K, Yada T. Resistin induces insulin resistance in pancreatic islets to impair glucose-induced insulin release. *Biochem Biophys Res Commun* 2007;353:1046-51.
74. Nakayamada S, Saito K, Umehara H, Ogawa N, Sumida T, Ito S, Minota S, Nara H, Kondo H, Okada J, Mimori T, Yoshifuji H, Sano H, Hashimoto N, Sugai S, Tanaka Y. Efficacy and safety of mizoribine for the treatment of Sjogren's syndrome: a multicenter open-label clinical trial. *Mod Rheumatol* 2007;17:464-9.
75. Namekawa M, Muriel MP, Janer A, Latouche M, Dauphin A, Debeir T, Martin E, Duyckaerts C, Prigent A, Depienne C, Sittler A, Brice A, Ruberg M. Mutations in the SPG3A gene encoding the GTPase atlastin interfere with vesicle trafficking in the ER/Golgi interface and Golgi morphogenesis. *Mol Cell Neurosci* 2007;35:1-13.
76. Naritomi H, Fujita T, Ito S, Ogihara T, Shimada K, Shimamoto K, Tanaka H,

- Yoshiike N. Design and baseline characteristics of an observational study in Japanese patients with hypertension: Japan Hypertension Evaluation with Angiotensin II Antagonist Losartan Therapy (J-HEALTH). *Hypertens Res* 2007;30:807-14.
77. Nonaka-Sarukawa M, Yamamoto K, Aoki H, Nishimura Y, Tomizawa H, Ichida M, Eizawa T, Muroi K, Ikeda U, Shimada K. Circulating endothelial progenitor cells in congestive heart failure. *Int J Cardiol* 2007;119:344-8.
78. Ogata H, Tokuyama K, Nagasaka S, Ando A, Kusaka I, Sato N, Goto A, Ishibashi S, Kiyono K, Struzik ZR, Yamamoto Y. Long-range correlated glucose fluctuations in diabetes. *Methods Inf Med* 2007;46:222-6.
79. Oh I, Ozaki K, Sato K, Meguro A, Tatara R, Hatanaka K, Nagai T, Muroi K, Ozawa K. Interferon-gamma and NF-kappaB mediate nitric oxide production by mesenchymal stromal cells. *Biochem Biophys Res Commun* 2007;355:956-62.
80. Oh I, Ozaki M, Miyazato A, Sato K, Meguro A, Muroi K, Nagai T, Mano H, Ozawa K. Screening of genes responsible for differentiation of mouse mesenchymal stromal cells by DNA micro-array analysis of C3H10T1/2 and C3H10T1/2-derived cell lines. *Cyotherapy* 2007;9:80-90.
81. Ohgushi M, Taniguchi A, Fukushima M, Nakai Y, Kuroe A, Ohya M, Nagasaka S, Taki Y, Yoshii S, Matsumoto K, Yamada Y, Inagaki N, Seino Y. Soluble tumor necrosis factor receptor 2 is independently associated with pulse wave velocity in nonobese Japanese patients with type 2 diabetes mellitus. *Metabolism* 2007;56:571-7.
82. Ohkuchi A, Hirashima C, Matsubara S, Suzuki H, Takahashi K, Arai F, Watanabe T, Kario K, Suzuki M. Alterations in placental growth factor levels before and after the onset of preeclampsia are more pronounced in women with early onset severe preeclampsia. *Hypertens Res* 2007;30:151-9.
83. Ohmiya N, Yano T, Yamamoto H, Arakawa D, Nakamura M, Honda W, Itoh A, Hirooka Y, Niwa Y, Maeda O, Ando T, Yao T, Matsui T, Iida M, Tanaka S, Chiba T, Sakamoto C, Sugano K, Goto H. Diagnosis and treatment of obscure GI bleeding at double balloon endoscopy. *Gastrointest Endosc* 2007;66:S72-7.
84. Oka S, Muroi K, Mori M, Matsuyama T, Fujiwara S, Oh I, Ono Y, Kikuchi S, Sato K, Ueda M, Toshima M, Ozaki K, Takatoku M, Nagai T, Ozawa K. Evaluation of platelet transfusion thresholds in patients with acute myeloblastic leukemia receiving induction chemotherapy. *Intern Med* 2007;46:1669-70.
85. Onishi A, Yamamoto H, Akimoto T, Saito O, Inoue M, Ando Y, Muto S, Kusano E. Reversible acute renal failure associated with clomipramine-induced interstitial nephritis. *Clin Exp Nephrol* 2007;11:241-3.
86. Otsubo S, Otsubo K, Sugimoto H, Ueda S, Otsubo Y, Otsubo O, Yajima A, Yagi S, Kataoka H, Iwasaki T, Iwasa Y, Takahashi M, Aoki A, Ino J, Uchida K, Yumura W, Akiba T, Nitta K. Characteristics of patients on hemodialysis therapy for more than 30 years. *Ther Apher Dial* 2007;11:274-9.
87. Ozaki K, Sato K, Oh I, Meguro A, Tatara R, Muroi K, Ozawa K. Mechanisms of immunomodulation by mesenchymal stem cells. *Int J Hematol* 2007;86:5-7.

88. Pickering TG, Eguchi K, Kario K. Masked hypertension: a review. *Hypertens Res* 2007;30:479-88.
89. Saito T, Saito O, Kawano T, Tamemoto H, Kusano E, Kawakami M, Ishikawa SE. Elevation of serum adiponectin and CD146 levels in diabetic nephropathy. *Diabetes Res Clin Pract* 2007;78:85-92.
90. Sakakura K, Ishikawa J, Okuno M, Shimada K, Kario K. Exaggerated ambulatory blood pressure variability is associated with cognitive dysfunction in the very elderly and quality of life in the younger elderly. *Am J Hypertens* 2007;20:720-7.
91. Sakamoto H, Mutoh H, Ido K, Satoh K, Hayakawa H, Sugano K. A close relationship between intestinal metaplasia and Cdx2 expression in human gallbladders with cholelithiasis. *Hum Pathol* 2007;38:66-71.
92. Sato K, Ozaki K, Oh I, Meguro A, Hatanaka K, Nagai T, Muroi K, Ozawa K. Nitric oxide plays a critical role in suppression of T-cell proliferation by mesenchymal stem cells. *Blood* 2007;109:228-34.
93. Sawada H, Hishida R, Hirata Y, Ono K, Suzuki H, Muramatsu S, Nakano I, Nagatsu T, Sawada M. Activated microglia affect the nigro-striatal dopamine neurons differently in neonatal and aged mice treated with 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine. *J Neurosci Res* 2007;85:1752-61.
94. Sekiya M, Yahagi N, Matsuzaka T, Takeuchi Y, Nakagawa Y, Takahashi H, Okazaki H, Iizuka Y, Ohashi K, Gotoda T, Ishibashi S, Nagai R, Yamazaki T, Kadokawa T, Yamada N, Osuga J, Shimano H. SREBP-1-independent regulation of lipogenic gene expression in adipocytes. *J Lipid Res* 2007;48:1581-91.
95. Seno H, Satoh K, Tsuji S, Shiratsuchi T, Harada Y, Hamajima N, Sugano K, Kawano S, Chiba T. Novel interleukin-4 and interleukin-1 receptor antagonist gene variations associated with non-cardia gastric cancer in Japan: comprehensive analysis of 207 polymorphisms of 11 cytokine genes. *J Gastroenterol Hepatol* 2007;22:729-37.
96. Shen WJ, Liang Y, Wang J, Harada K, Patel S, Michie SA, Osuga J, Ishibashi S, Kraemer FB. Regulation of hormone-sensitive lipase in islets. *Diabetes Res Clin Pract* 2007;75:14-26.
97. Shiizaki K, Hayakawa N, Imazeki I, Hatamura I, Okada T, Negi S, Sakaguchi T, Shigematsu T, Akizawa T. Binding of highly concentrated maxacalcitol to the nuclear vitamin D receptors of parathyroid cells. *Nephrol Dial Transplant* 2007;22:1078-86.
98. Shimazaki H, Ando Y, Nakano I, Dalmau J. Reversible limbic encephalitis with antibodies against the membranes of neurones of the hippocampus. *J Neurol Neurosurg Psychiatry* 2007;78:324-5.
99. Shimazaki H, Sakoe K, Niijima K, Nakano I, Takiyama Y. An unusual case of a spasticity-lacking phenotype with a novel SACS mutation. *J Neurol Sci* 2007;255:87-9.
100. Shimizu T, Murata M, Tomizawa H, Mitsuhashi T, Katsuki T, Shimada K. Systemic lupus erythematosus initially manifesting as acute pericarditis complicating with cardiac tamponade : a case report. *J Cardiol* 2007;49:273-6.
101. Soda M, Choi YL, Enomoto M, Takada S, Yamashita Y, Ishikawa S, Fujiwara S,

- Watanabe H, Kurashina K, Hatanaka H, Bando M, Ohno S, Ishikawa Y, Aburatani H, Niki T, Sohara Y, Sugiyama Y, Mano H. Identification of the transforming EML4-ALK fusion gene in non-small-cell lung cancer. *Nature* 2007;448:561-6.
102. Sone H, Yoshimura Y, Tanaka S, Iimuro S, Ohashi Y, Ito H, Seino H, Ishibashi S, Akanuma Y, Yamada N. Cross-sectional association between BMI, glycemic control and energy intake in Japanese patients with type 2 diabetes. Analysis from the Japan Diabetes Complications Study. *Diabetes Res Clin Pract* 2007;77 Suppl 1:S23-9.
103. Stevens KR, Rolle MW, Minami E, Ueno S, Nourse MB, Virag JI, Reinecke H, Murry CE. Chemical dimerization of fibroblast growth factor receptor-1 induces myoblast proliferation, increases intracardiac graft size, and reduces ventricular dilation in infarcted hearts. *Hum Gene Ther* 2007;18:401-12.
104. Sugano K. Prevention of gastric cancer: urgent need to implement Helicobacter pylori eradication therapy as a primary preventive measure in Japan. *J Gastroenterol* 2007;42 Suppl 17:1-2.
105. Sugano K, Marcon N. The First International Workshop on Double Balloon Endoscopy: a consensus meeting report. *Gastrointest Endosc* 2007;66:S7-11.
106. Sunada K, Yamamoto H, Hayashi Y, Sugano K. Clinical importance of the location of lesions with regard to mesenteric or antimesenteric side of the small intestine. *Gastrointest Endosc* 2007;66:S34-8.
107. Tajima S, Bando M, Ohno S, Sugiyama Y, Oshikawa K, Tominaga S, Itoh K, Takada T, Suzuki E, Gejyo F. ST2 gene induced by type 2 helper T cell (Th2) and proinflammatory cytokine stimuli may modulate lung injury and fibrosis. *Exp Lung Res* 2007;33:81-97.
108. Tajima S, Bando M, Yamasawa H, Ohno S, Moriyama H, Terada M, Takada T, Suzuki E, Gejyo F, Sugiyama Y. Preventive effect of hochu-ekki-to, a Japanese herbal medicine, on bleomycin-induced lung injury in mice. *Respirology* 2007;12:814-22.
109. Takahashi M, Otsubo S, Takei T, Sugiura H, Yoshida K, Tamei N, Koike M, Uchida K, Yumura W, Kawamura S, Horita S, Akiba T, Nitta K. Anti-glomerular basement membrane antibody disease with granulomatous lesions on renal biopsy. *Intern Med* 2007;46:295-301.
110. Takahashi M, Otsubo S, Uchida K, Yumura W, Nitta K. Association between serum adiponectin levels and arteriolosclerosis in IgA nephropathy patients. *Intern Med* 2007;46:453-9.
111. Takahashi M, Tanaka T, Azuma M, Kusano E, Aikawa T, Shibayama T, Yazaki Y, Mizuo H, Inoue J, Okamoto H. Prolonged fecal shedding of hepatitis E virus (HEV) during sporadic acute hepatitis E: evaluation of infectivity of HEV in fecal specimens in a cell culture system. *J Clin Microbiol* 2007;45:3671-9.
112. Takatoku M, Uchiyama T, Okamoto S, Kanakura Y, Sawada K, Tomonaga M, Nakao S, Nakahata T, Harada M, Murate T, Ozawa K. Retrospective nationwide survey of Japanese patients with transfusion-dependent MDS and aplastic anemia highlights the negative impact of iron overload on morbidity/mortality. *Eur J Haematol* 2007;78:487-94.
113. Takei T, Koike M, Suzuki K, Shirota S, Itabashi M, Ohtsubo S, Sugiura H, Kojima C, Takahashi M, Ino J, Ogawa T, Uchida K, Tsuchiya K, Yumura W, Nitta K. The characteristics

of relapse in adult-onset minimal-change nephrotic syndrome. *Clin Exp Nephrol* 2007;11:214-7.

114. Takei Y, Mizukami H, Saga Y, Yoshimura I, Hasumi Y, Takayama T, Kohno T, Matsushita T, Okada T, Kume A, Suzuki M, Ozawa K. Suppression of ovarian cancer by muscle-mediated expression of soluble VEGFR-1/Flt-1 using adeno-associated virus serotype 1-derived vector. *Int J Cancer* 2007;120:278-84.
115. Takiyama Y. Sacsinopathies: sacsin-related ataxia. *Cerebellum* 2007;1-7.
116. Tamada K, Wada S, Nakazawa K, Hatanaka H, Tomiyama T, Ohashi A, Sugano K, Ido K. Teaching deep cannulation of the bile duct during endoscopic retrograde cholangiography. *Digestive Endoscopy* 2007;9:195-200.
117. Tanaka M, Yamazaki S, Hayashino Y, Fukuwara S, Akiba T, Saito A, Asano Y, Port FK, Kurokawa K, Akizawa T. Hypercalcaemia is associated with poor mental health in haemodialysis patients: results from Japan DOPPS. *Nephrol Dial Transplant* 2007;22:1658-64.
118. Tanaka T, Takahashi M, Kusano E, Okamoto H. Development and evaluation of an efficient cell-culture system for Hepatitis E virus. *J Gen Virol* 2007;88:903-11.
119. Taniguchi A, Fukushima M, Kuroe A, Sakaguchi K, Hashimoto H, Yoshioka I, Kitatani N, Tsuji T, Ohya M, Ohgushi M, Nagasaka S, Isogai O, Nakai Y, Inagaki N, Seino Y. Metabolic syndrome, insulin resistance, and atherosclerosis in Japanese type 2 diabetic patients. *Metabolism* 2007;56:1099-103.
120. Tarumoto T, Imagawa S, Kobayashi M, Hirayama A, Ozawa K, Nagasawa T. L-arginine administration reverses anemia associated with renal disease. *Int J Hematol* 2007;86:126-9.
121. Thompson AM, Eguchi K, Reznik ME, Shah SS, Pickering TG. Validation of an oscillometric home blood pressure monitor in an end-stage renal disease population and the effect of arterial stiffness on its accuracy. *Blood Press Monit* 2007;12:227-32.
122. Tomizawa H, Yamamoto K, Iwata T, Nishimura Y, Nomura Y, Hoshida Y, Eguchi K, Kaminishi Y, Misawa Y, Hironaka M, Shimada K. [Papillary fibroelastoma in the right ventricular outflow tract: a case report]. *J Cardiol* 2007;49:199-203.
123. Uchida M, Kiritó K, Endo H, Ozawa K, Komatsu N. Activation of FKHLR1 plays an important role in protecting erythroid cells from erythropoietin deprivation-induced apoptosis in a human erythropoietin-dependent leukemia cell line, UT-7/EPO. *Int J Hematol* 2007;86:315-24.
124. Ueno S, Weidinger G, Osugi T, Kohn AD, Golob JL, Pabon L, Reinecke H, Moon RT, Murry CE. Biphasic role for Wnt/beta-catenin signaling in cardiac specification in zebrafish and embryonic stem cells. *Proc Natl Acad Sci U S A* 2007;104:9685-90.
125. Usui R, Shibuya M, Ishibashi S, Maru Y. Ligand-independent activation of vascular endothelial growth factor receptor 1 by low-density lipoprotein. *EMBO Rep* 2007;8:1155-61.
126. Yamamoto H. Foreword: double balloon endoscopy. *Gastrointest Endosc* 2007;66:S2.
127. Yamamoto H. Technology insight: endoscopic submucosal dissection of gastrointestinal neoplasms. *Nat Clin Pract Gastroenterol Hepatol* 2007;4:511-20.

128. Yamamoto K. Synchronicity and ventricular arrhythmias in hypertension. *Hypertens Res* 2007;30:755-6.
129. Yamashita Y, Minoura K, Taya T, Fujiwara SI, Kurashina K, Watanabe H, Choi YL, Soda M, Hatanaka H, Enomoto M, Takada S, Mano H. Analysis of chromosome copy number in leukemic cells by different microarray platforms. *Leukemia* 2007;21:1333-7.
130. Yamazaki M, Takei T, Otsubo S, Iwasa Y, Yabuki Y, Suzuki K, Koike M, Uchida K, Tsuchiya K, Yumura W, Horita S, Honda K, Akiba T, Nitta K. Wegener's granulomatosis complicated by intestinal ulcer due to cytomegalovirus infection and by thrombotic thrombocytopenic purpura. *Intern Med* 2007;46:1435-40.
131. Yamazaki T, Goto S, Shigematsu H, Shimada K, Uchiyama S, Nagai R, Yamada N, Matsumoto M, Origasa H, Bhatt DL, Steg PG, Ikeda Y. Prevalence, awareness and treatment of cardiovascular risk factors in patients at high risk of atherothrombosis in Japan. *Circ J* 2007;71:995-1003.
132. Yanagawa T, Taniguchi A, Fukushima M, Nakai Y, Nagasaka S, Ohgushi M, Matsumoto K, Kuroe A, Ohya M, Seino Y. Leptin, triglycerides, and interleukin 6 are independently associated with C-reactive protein in Japanese type 2 diabetic patients. *Diabetes Res Clin Pract* 2007;75:2-6.
133. Yano Y, Hoshide S, Ishikawa J, Hashimoto T, Eguchi K, Shimada K, Kario K. Differential impacts of adiponectin on low-grade albuminuria between obese and nonobese persons without diabetes. *J Clin Hypertens (Greenwich)* 2007;9:775-82.
134. Yano Y, Hoshide S, Ishikawa J, Noguchi C, Tukui D, Takanori H, Tada M, Kanemaru Y, Yano A, Ishikawa S, Shimada K, Kario K. The differential effects of angiotensin II type 1 receptor blockers on microalbuminuria in relation to low-grade inflammation in metabolic hypertensive patients. *Am J Hypertens* 2007;20:565-72.
135. Yasuda H, Hirata S, Inoue K, Mashima H, Ohnishi H, Yoshioka M. Involvement of membrane-type bile acid receptor M-BAR/TGR5 in bile acid-induced activation of epidermal growth factor receptor and mitogen-activated protein kinases in gastric carcinoma cells. *Biochem Biophys Res Commun* 2007;354:154-9.
136. Yokoyama M, Origasa H, Matsuzaki M, Matsuzawa Y, Saito Y, Ishikawa Y, Oikawa S, Sasaki J, Hishida H, Itakura H, Kita T, Kitabatake A, Nakaya N, Sakata T, Shimada K, Shirato K. Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomised open-label, blinded endpoint analysis. *Lancet* 2007;369:1090-8.
137. Yoshio T, Okamoto H, Minota S. Antibodies to bovine serum albumin do not affect the results of enzyme-linked immunosorbent assays for IgG anti-NR2 glutamate receptor antibodies: reply to the letter by Hirohata et al. *Arthritis Rheum* 2007;56:2813-4.