

論文（主な英文論文のみ）

1. Primadharsini PP, Nagashima S, Takahashi M, Kobayashi T, Nishiyama T, Nishizawa T, Yasuda J, Mulyanto, Okamoto H. Multivesicular body sorting and the exosomal pathway are required for the release of rat hepatitis E virus from infected cells. *Virus Res.* 2020;278:197868
2. Nishiyama T, Kobayashi T, Jirintai S, Nagashima S, Primadharsini PP, Nishizawa T, Okamoto H. Antiviral candidates against the hepatitis E virus (HEV) and their combinations inhibit HEV growth in in vitro. *Antiviral Res.* 2019;170:104570
3. Nishiyama T, Kobayashi T, Jirintai S, Kii I, Nagashima S, Prathiwi Primadharsini P, Nishizawa T, Okamoto H. Screening of novel drugs for inhibiting hepatitis E virus replication. *J Virol Methods* 2019;270:1-11
4. Tanggis, Kobayashi T, Takahashi M, Jirintai S, Nishizawa T, Nagashima S, Nishiyama T, Kunita S, Hayama E, Tanaka T, Mulyanto, Okamoto H. An analysis of two open reading frames (ORF3 and ORF4) of rat hepatitis E virus genome using its infectious cDNA clones with mutations in ORF3 or ORF4. *Virus Res* 2018;249:16-30
5. Primadharsini PP, Miyake M, Kunita S, Nishizawa T, Takahashi M, Nagashima S, Tanggis, Ohnishi H, Kobayashi T, Nishiyama T, Jirintai S, Okamoto H. Full-length genome of a novel genotype 3 hepatitis E virus strain obtained from domestic pigs in Japan. *Virus Res* 2017;240:147-153
6. Nagashima S, Takahashi M, Kobayashi T, Tanggis, Nishizawa T, Nishiyama T, Primadharsini PP, Okamoto H. The characterization of the quasi-enveloped hepatitis E virus particles released by the cellular exosomal pathway. *J Virol* 2017;91(22):e00822-17
7. Kobayashi T, Takahashi M, Tanggis, Mulyanto, Jirintai S, Nagashima S, Nishizawa T, Okamoto H. Characterization and epitope mapping of monoclonal antibodies raised against rat hepatitis E virus capsid protein: An evaluation of their neutralizing activity in a cell culture system. *J Virol Methods* 2016;233:78-88
8. Nishizawa T, Hoshino T, Naganuma A, Kobayashi T, Nagashima S, Takahashi M, Takagi H, Okamoto H. Enhanced pregenomic RNA levels and lowered precore mRNA transcription efficiency in a genotype A hepatitis B virus genome with C1766T and T1768A mutations obtained from a fulminant hepatitis patient. *J Gen Virol* 2016;97(10):2643-2656
9. Takahashi M, Kobayashi T, Tanggis, Jirintai S, Mulyanto, Nagashima S, Nishizawa T, Kunita S, Okamoto H. Production of monoclonal antibodies against the ORF3 protein of rat hepatitis E virus (HEV) and demonstration of the incorporation of the ORF3 protein into enveloped rat HEV particles. *Arch Virol* 2016;161(12):3391-3404

10. Nagashima S, Kobayashi T, Tanaka T, Tanggis, Jirintai S, Takahashi M, Nishizawa T, Okamoto H. Analysis of adaptive mutations selected during the consecutive passages of hepatitis E virus produced from an infectious cDNA clone. *Virus Res* 2016;223:170-180
11. Tsatsralt-Od B, Baasanjav N, Nyamkhuu D, Ohnishi H, Takahashi M, Kobayashi T, Nagashima S, Nishizawa T, Okamoto H. Molecular analysis of hepatitis A virus strains obtained from patients with acute hepatitis A in Mongolia, 2004-2013. *J Med Virol* 2016;88(4):622-630
12. Kobayashi T, Machida K, Imataka H. Human cell extract-derived cell-free systems for virus synthesis. *Methods Mol Biol* 2014;1118:149-156
13. Nagashima S, Jirintai S, Takahashi M, Kobayashi T, Tanggis, Nishizawa T, Kouki T, Yashiro T, Okamoto H. Hepatitis E virus egress depends on the exosomal pathway, with secretory exosomes derived from multivesicular bodies. *J Gen Virol* 2014;95(Pt 10):2166-2175
14. Nagashima S, Takahashi M, Jirintai S, Tanggis, Kobayashi T, Nishizawa T, Okamoto H. The membrane on the surface of hepatitis E virus particles is derived from the intracellular membrane and contains trans-Golgi network protein 2. *Arch Virol* 2014;159(5):979-991
15. Jirintai S, Tanggis, Mulyanto, Suparyatmo JB, Takahashi M, Kobayashi T, Nagashima S, Nishizawa T, Okamoto H. Rat hepatitis E virus derived from wild rats (*Rattus rattus*) propagates efficiently in human hepatoma cell lines. *Virus Res* 2014;185:92-102
16. Machida K, Mikami S, Masutani M, Mishima K, Kobayashi T, Imataka H. A Translation system reconstituted with human factors proves that processing encephalomyocarditisvirus proteins 2A and 2B occurs in the elongation phase of translation without eukaryotic release factors. *J Biol Chem* 2014;289(46):31960-31971
17. Kobayashi T, Yukigai J, Ueda K, Machida K, Masutani M, Nishino Y, Miyazawa A, Imataka H. Purification and visualization of encephalomyocarditisvirus synthesized by an in vitro protein expression system derived from mammalian cell extract. *Biotechnol Lett* 2013;35(3):309-314
18. Masutani M, Machida K, Kobayashi T, Yokoyama S, Imataka H. Reconstitution of eukaryotic translation initiation factor 3 by co-expression of the subunits in a human cell-derived in vitro protein synthesis system. *Protein Expr Purif* 2013;87(1):5-10
19. Kobayashi T, Nakamura Y, Mikami S, Masutani M, Machida K, Imataka H. Synthesis of encephalomyocarditis virus in a cell-free system: from DNA to RNA virus in one tube. *Biotechnol Lett* 2012;34(1):67-73
20. Jirintai S, Jinshan, Tanggis, Manglai D, Mulyanto, Takahashi M, Nagashima S, Kobayashi T, Nishizawa T, Okamoto H. Molecular analysis of hepatitis E virus from farm rabbits in Inner Mongolia, China and its successful propagation in A549 and PLC/PRF/5 cells. *Virus Res*

2012;170(1-2):126-137

21. Kobayashi T, Machida K, Mikami S, Masutani M, Imataka H. Cell-free RNA replication systems based on a human cell extracts-derived in vitro translation system with the encephalomyocarditisvirus RNA. J Biochem 2012;150(4):423-430

※日本生化学会 JB 論文賞受賞(2012)

22. Mikami S, Kobayashi T, Machida K, Masutani M, Yokoyama S, Imataka H. N-terminally truncated GADD34 proteins are convenient translation enhancers in a human cell-derived in vitro protein synthesis system. Biotechnol Lett 2010;32(7):897-902

23. Mikami S, Kobayashi T, Imataka H. Cell-free protein synthesis systems with extracts from cultured human cells. Methods Mol Biol 2010;607:43-52

24. Mikami S, Kobayashi T, Masutani M, Yokoyama S, Imataka H. A human cell-derived in vitro coupled transcription/translation system optimized for production of recombinant proteins. Protein Expr Purif 2008;62(2):190-198

25. Kobayashi T, Mikami S, Yokoyama S, Imataka H. An improved cell-free system for picornavirus synthesis. J Virol Methods 2007;142(1-2):182-188

26. Mikami S, Kobayashi T, Yokoyama S, Imataka H. A hybridoma-based in vitro translation system that efficiently synthesizes glycoproteins. J Biotechnol 2006;127(1):65-78

27. Yoshida S, Kobayashi T, Matsuoka H, Seki C, Gosnell WL, Chang SP, Ishii A. T-cell activation and cytokine production via a bispecific single-chain antibody fragment targeted to blood-stage malaria parasites. Blood 2003;101(6):2300-2306